

Unitile: a finite, decisive tile-sliding game

Mark Steere

October 2026

Abstract

Unitile is a game for two players on a 5×5 board. Every cell but one holds a red tile or a blue tile. The one unoccupied cell is called the *bubble*. A move slides a tile into the bubble from an orthogonally adjacent cell. Red tries to unite all the red tiles into one orthogonally connected group. Blue tries to unite all the blue tiles. When a move unites a color, the player of that color wins. Without a further rule, the players could slide tiles indefinitely, never reaching a decisive conclusion. Unitile provides a rule that keeps play marching steadily toward a victory. A cycle or a deadlocked position can never occur. Unitile is naturally finite. No separate, tacked-on rule expressly prohibiting repeated positions is required. Applying the rule requires a computer program. The rule uses a function that gives each color a value, the average number of random slides needed to unite the tiles of that color, starting from the current position. Each value is an exact number, not an estimate. Every move must lower the total of the two values, except a move that unites a color. Such a move ends the game and is always legal. We prove that the player to move always has at least one legal move. A computer confirmed the same fact in every position on the board. We also show that a simpler rule of the same kind, but based on the fewest slides needed to unite each color, leaves some positions with no legal move.

1 Introduction

Every cell of the Unitile board but one holds a red tile or a blue tile. The one unoccupied cell is called the *bubble*. On your turn, you slide a tile of either color into the bubble from an orthogonally adjacent cell. The cell that the tile leaves becomes the new bubble. Red's goal is to unite all the red tiles into a single connected group. Blue's goal is to do the same with the blue tiles.

As the game has been described so far in this introduction, players could shuffle the tiles indefinitely, never reaching a conclusive outcome. A position could repeat after a number of moves, or you could simply slide the last moved tile back to where it came from. The game needs one more rule, a rule that makes the game decisive. The rule must guarantee that the player to move always has a legal move and that someone wins.

You might try a rule that forbids repeating a position, a rule known as superko. Superko has three problems in Unitile: 1) a position could arise in which every possible move would repeat an earlier position, leaving the player to move with no legal move, 2) a move could suddenly be forbidden because it repeats a position that both players have long since forgotten, and 3) superko is an ugly rule, bolted onto the game after the fact. The first problem is common. In 20,000 games in which each player chose at random among the moves that superko allows, 30% of the games deadlocked in a position in which the player to move had no legal move.

What is needed is a rule under which every move brings the game closer to a decisive conclusion and the player to move always has a legal move. Unitile uses a function for this purpose. In every position, the function gives each color a value that measures how far the tiles of that color are from

being united. Every move must lower the total of the two values, except a move that ends the game by uniting a color.

The challenge was to find a function that never leaves a player without a legal move. The player to move is left without a legal move when no move unites a color and every possible move hurts one color at least as much as it helps the other color. Obvious choices of function all fail somewhere. Here are two ways in which an obvious function can fail.

1) In some positions, every possible move breaks up a group of one color without bringing the other color any closer to being united. For example, if the function counts the groups of each color, every move raises the total of the two values.

2) In other positions, the two colors need the bubble in different places. Every possible move brings the bubble closer to where one color needs it and farther from where the other color needs it. If the function counts the slides that each color still needs, every such move helps one color by one slide and hurts the other color by one slide. No move lowers the total.

One function that *does* work is built from random sliding. For each color, imagine sliding tiles at random until the tiles of that color happen to unite. The average number of slides needed is the color's value, an exact number. We prove that at least one move always lowers the total of the two values. Determining a color's value requires an extensive calculation that cannot be worked out by hand. A computer must be used to calculate the values.

2 The rules

Unitile is played on a 5×5 board of cells. Every cell but one holds a tile. Each tile is red or blue. The board has twelve red tiles and twelve blue tiles. The one unoccupied cell is called the *bubble*.

In the starting position, the bubble is the center cell. The 24 tiles fill the other cells in reading order, from left to right within each row, and row by row from the top. The colors of the tiles alternate in that order. The first tile, in the top left corner, is blue. Figure 1 shows the starting position.




Figure 1: The starting position. The bubble is the unoccupied cell in the center.

The players alternate sliding tiles, one slide per turn, starting with Red. The player to move chooses a tile of either color that is orthogonally adjacent to the bubble and slides it into the bubble. The cell that the tile has left becomes the new bubble. Figure 2 shows a move.




Figure 2: A move. In (a), the red tile above the bubble slides down into it. In (b), the cell that the red tile left is the new bubble.

A *group* is a maximal set of orthogonally interconnected, same-colored tiles. A color is *united* when all of its tiles form a single group.

The game ends as soon as a move unites a color. The player of that color wins. A move can unite the opponent's color, and a player who makes such a move loses. No move unites both colors at once, because a move changes the cell of only one tile.

These rules alone do not make the game end. For example, after Red slides a tile into the bubble, Blue may slide the same tile back, and the position is exactly the same as before. One further rule decides which moves are legal. Before stating the rule, we describe what it must achieve.

3 What a progress rule must do

We look for a function that gives each color a value in every position. A position here means the arrangement of the tiles and the bubble on the board. The value of a color measures how far the tiles of that color are from being united. Two requirements shape the function:

1) Every move must lower the total of the two values, except a move that unites a color and ends the game. As long as the game goes on, the total can never equal or exceed an earlier total. No position can repeat.

2) The player to move must have at least one legal move in every position. A position in which the player to move has no legal move is called a *deadlock*. We ask for no deadlocks in any position.

Together, the two requirements bring the game closer to a decisive conclusion with every move.

4 A function that fails

Before giving the rule, we look at a natural attempt at a rule and at what goes wrong with it. This natural attempt uses the fewest slides required to unite a color as the value of the color.

In the starting position, each color needs 16 slides. Consider the rule that every move must lower the total of the fewest slides for red and the fewest slides for blue, unless the move unites a color. Under this rule, only four different games can be played from the starting position. Every one of the four games reaches a deadlock.

Figure 3(a) shows the position in which one of the four games reaches a deadlock, after five moves. In this position, red needs 12 slides and blue needs 11. The bubble has three neighbors. Sliding the blue tile above the bubble down gives the position of Figure 3(b), in which red needs 13 slides and blue needs 12. In Figure 3(a), sliding the red tile below the bubble up gives the position of Figure 3(c), in which red needs 11 slides and blue needs 12. In Figure 3(a), sliding the red tile to the right of the bubble to the left gives the position of Figure 3(d), in which red needs 13 slides and blue needs 10. No move unites a color or lowers the total of 23. The position is a deadlock.




Figure 3: A game under the rule built on the fewest slides can reach the position in (a), five moves from the starting position. The three possible moves in (a) lead to (b), (c), and (d). Each of the five moves that lead to (a) lowers the total of the fewest slides. No move from (a) lowers the total.

5 Random-slide values

5.1 The definition

Imagine sliding tiles at random. At each step, one of the tiles orthogonally adjacent to the bubble is chosen at random, each with the same chance. The chosen tile slides into the bubble. The bubble has two, three, or four neighbors, depending on where it is. Whether the chosen tile is red or blue makes no difference to the choice.

Definition 1. Let P be a position. Red's value $r(P)$ is the average number of random slides, starting from P , until the red tiles are united. Blue's value $b(P)$ is the average number of random slides, starting from P , until the blue tiles are united. When the red tiles are united in P , $r(P) = 0$. When the blue tiles are united in P , $b(P) = 0$.

The random sliding behind $r(P)$ goes on until the red tiles are united, even if the blue tiles become united first.

The random sliding unites the red tiles sooner or later. From every position, some sequence of slides unites the red tiles. From wherever the random sliding happens to be, random sliding follows such a sequence with a chance that is small but never zero. The chance that random sliding goes on forever without uniting the red tiles is zero, and $r(P)$ is a finite average. The same holds for $b(P)$.

An average of a random quantity can be an exact number. The roll of a die, for example, is random, but the average roll is exactly 3.5. In the same way, $r(P)$ is one exact number for each position P .

5.2 The Unitile rule

Definition 2. The player to move must make a move that either unites a color or lowers the total $r(P) + b(P)$.

A move that unites a color ends the game. The rule always allows such a move, even a move that unites the opponent's color or does not lower the total. Every other legal move lowers the total.

The Unitile rule meets the first requirement of a progress rule. Every legal move either ends the game or lowers the total. The Unitile rule also meets the second requirement. No position is a deadlock. The proof comes next.

5.3 The averaging equation

Lemma 3. *Let P be a position in which the red tiles are not united. Let $P_1, \dots, P_k$ be the positions that the k possible slides lead to, where k is 2, 3, or 4. Then*

$$r(P) = 1 + \frac{r(P_1) + \dots + r(P_k)}{k}.$$

The same equation holds for Blue's value, in every position in which the blue tiles are not united.


Proof. One random slide from P leads to $P_1, P_2, \dots$, or P_k , each with the same chance $1/k$. If the first slide leads to P_i , random sliding needs, on average, $r(P_i)$ further slides, totaling $1 + r(P_i)$ slides. Red's value $r(P)$ is the average of these k averages.

$$\begin{aligned} r(P) &= \frac{(1 + r(P_1)) + (1 + r(P_2)) + \dots + (1 + r(P_k))}{k} \\ &= \frac{k + r(P_1) + r(P_2) + \dots + r(P_k)}{k} \\ &= \frac{k}{k} + \frac{r(P_1) + r(P_2) + \dots + r(P_k)}{k} \\ &= 1 + \frac{r(P_1) + r(P_2) + \dots + r(P_k)}{k}. \end{aligned}$$

□


The averaging equation is a standard result of probability theory [1, Section 1.3]. In probability theory, the average number of random steps needed to reach a goal is called an expected hitting time. As far as we know, expected hitting times have not been used before to build the rule of a game.

In words, the averaging equation says that the average of Red's values one slide after P is exactly 1 less than Red's value in P . Figure 4 shows the values in the starting position and after each of the four possible first moves. Both colors start with the value 1147.1914. After the four first moves, Red's values are 1146.4013, 1145.6054, 1148.5855, and 1144.1736. The average of these four values is 1146.1914, exactly 1 less than 1147.1914. Blue's values after the four first moves are the same four values in a different order, and their average is again exactly 1 less than Blue's value in the starting position.




Red 1147.1914
Blue 1147.1914
total 2294.3829

(a)




Red 1146.4013
Blue 1144.1736
total 2290.5749

(b)




Red 1145.6054
Blue 1148.5855
total 2294.1908

(c)



Red 1148.5855
Blue 1145.6054
total 2294.1908

(d)



Red 1144.1736
Blue 1146.4013
total 2290.5749

(e)

Figure 4: The random-slide values in the starting position (a) and after each of the four possible first moves (b), (c), (d), and (e). All four first moves lower the total, all four first moves are legal, and the four totals after the first moves average exactly 2 less than the total in the starting position.

5.4 No deadlocks

Theorem 4. *In every position in which neither color is united, at least one move lowers the total $r(P) + b(P)$ by 2 or more.*

Proof. Let P be a position in which neither color is united. Let $P_1, \dots, P_k$ be the positions that the k possible moves lead to. The red tiles are not united in P . By the averaging equation for Red, the average of $r(P_1), \dots, r(P_k)$ is $r(P) - 1$. The blue tiles are not united in P either. By the averaging equation for Blue, the average of $b(P_1), \dots, b(P_k)$ is $b(P) - 1$. The average of the totals $r(P_i) + b(P_i)$ is the sum of the two averages, $r(P) + b(P) - 2$. In any average, at least one of the numbers being averaged is at most the average. At least one of the totals $r(P_i) + b(P_i)$ is at most $r(P) + b(P) - 2$, and the move to that position lowers the total by 2 or more. $\square$

Corollary 5. *Under the Unitile rule, no position is a deadlock, and every game ends in a win for one of the players.*

Proof. By Theorem 4, every position in which neither color is united has a legal move, and no position is a deadlock. Every legal move either unites a color, which ends the game, or lowers the total. As long as the game goes on, the total keeps dropping, and no position can occur twice. The board has finitely many positions. A game that went on forever would have to repeat a position. Every game must end. The last move of every game unites a color and decides the winner. $\square$

In the starting position, the total is 2294.3829. Two of the four first moves lower the total by 3.81, and the other two lower it by 0.19. The four totals after the first moves average 2292.3829,

exactly 2 less than the starting total.

Figure 5 shows the position after the first move of Figure 4(b), in which the red tile above the bubble slid down. The bubble again has four neighbors. Three of the four moves lower the total and are legal. The fourth move slides the red tile back up. That move returns to the starting position, raises the total, and is not legal. Red's values in panels (b), (c), (d), and (e) average exactly 1 less than Red's value in panel (a), and the same holds for Blue's values.




Figure 5: One move later. In (a), from the starting position, the red tile above the bubble has slid down. The moves of (b), (c), and (d) lower the total and are legal. The move of (e) slides the red tile back up, and the position of (e) is again the starting position. The move of (e) raises the total and is not legal.

The random-slide function succeeds where the fewest-slides function failed. In about 40% of all slides, one color's fewest-slides value rises by 1, the other color's fewest-slides value falls by 1, and the total stays the same. The random-slide values change by uneven amounts. In Figure 5(c), Blue's value drops by more than 4.6, and Red's value rises by less than 1.7. The averaging equation guarantees that, in every position, one color's drop outweighs the other color's rise for at least one move.

5.5 Computing the values

A computer finds the values by solving all of the averaging equations together, not by sliding tiles at random. The values are computed once, in advance, and stored in a table. Every game uses the same table.

As a check, the computer also tested Theorem 4 position by position. The board has 64,087,986 positions in which neither color is united. Every one of these positions has at least one legal move. The programs used for this paper are available from the author.

6 Discussion

The proof of Theorem 4 uses nothing special about the 5×5 board. The proof works on any board, with any numbers of red and blue tiles, on which each color can be united from every position.

Unitile is played on the 5×5 board for other reasons. A square board with an even number of rows leaves an odd number of cells for the tiles. On such a board, the two colors cannot have the same number of tiles. The 3×3 board is too small for an interesting game. On the 7×7 board, there are about 1.6×10^{15} positions. Computing their values is far beyond the computer used for this paper.

The values cannot be worked out by hand. A program can show the players which moves are legal. The players choose among those moves.

An open question remains. Is there a function that a player can compute in his head and that still leaves no deadlocks?

Acknowledgment

The analysis, the proofs, and the computer checks of this paper were produced by Claude, an AI model made by Anthropic. The text was written jointly by Mark Steere and Claude.

References

- [1] J. R. Norris, *Markov Chains*, Cambridge University Press, Cambridge, 1997.